



FORMERLY WILLOW RUN LABORATORIES, THE UNIVERSITY OF MICHIGAN

P. O. BOX 618 • ANN ARBOR • MICHIGAN • 48107

PHONE (313) 483-0500

E7.4-10394

CR-137212

101800-10-L
20 March 1974

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Oil Pollution Detection, Monitoring and Law Enforcement
Quarterly Progress Report, February 1974

(E74-10394) OIL POLLUTION DETECTION,
MONITORING AND LAW ENFORCEMENT Quarterly
Progress Report (Environmental Research
Inst. of Michigan) 2 p HC \$4.00

N74-19003

Unclas
00394

CSCI 08J G3/13

EREP Investigation 417
NASA Contract NAS9-13281

Prepared by

Robert Horvath - Principal Investigator

NASA Technical Monitor

Mr. Timothy White/TF6
National Aeronautics Space Administration
Johnson Space Center
Principal Investigator Management Office
Houston, Texas 77058



101800-10-L


Oil Pollution Detection, Monitoring and Law Enforcement
Quarterly Progress Report, February 1974

This report covers progress during the fourth quarter of Contract NAS9-13281, "Evaluate Skylab EREP Data for Oil Pollution Detection, Monitoring and Law Enforcement", EREP No. 417. The work is being conducted in the Infrared and Optics Division of the Environmental Research Institute of Michigan, under the general supervision of Mr. R. R. Legault. The principal investigator is Mr. R. Horvath.


An extensive review of U. S. Coast Guard and EPA pollution report files was undertaken near the end of the report period. The purpose of this review was to identify whether any significant oil slicks were present in areas and at times coincident with EREP data takes. It was found that no confirmed coincidences exist. Thus, it now appears that we may not be able to obtain EREP data from known and documented oil slicks for analysis.

In the absence of documented slicks, a possible alternative approach is to attempt to locate unreported slicks by interpretation of available EREP data. This could be attempted by utilizing available data from areas wherein a high probability for slicks exist. For example, areas of natural oil seepage in the Gulf of Mexico and along the coast of southern California would be appropriate. This approach will be evaluated during the next quarterly period.

Respectfully submitted,


Robert Horvath
Principal Investigator

APPROVED BY:


Richard R. Legault
Director, Infrared and Optics Division

RH/RRL/dac